

Pacific Wildcat Resources Corp.: Further Wide and High Grade Rare Earth Intercepts at Mrima Hill Confirm High Grade Rare Earth Mineralised Zone

15.07.2013 | [Marketwired](#)

VANCOUVER, BRITISH COLUMBIA--(Marketwired - Jul 15, 2013) - [Pacific Wildcat Resources Corp.](#) (TSX VENTURE:PAW) ("**PAW**" or the "**Company**") is pleased to report additional assay results following on from those reported in its News Release dated 26th June 2013 (See NR 2013-11 titled "**Wide and High Grade Rare Earth intercepts show Mrima Hill Mineralisation Open to the South**") from the next 655 metres (7 holes) of Reverse Circulation ("**RC**") drilling at the Mrima Hill Niobium and Rare Earth Project (PAW indirect 70% ownership) in Kenya.

These assay results are from holes drilled across the centre of what is historically understood to be a higher grade zone of Rare Earth mineralization at Mrima Hill. Results continue to demonstrate the presence of Wide and High Grade zones of mineralization and remain open laterally and at depth. In addition elevated niobium ("**Nb₂O₅**") results have also been intersected in a number of holes further confirming the potential of this large mineralized system.

Highlights include:

Rare Earths

- Wide zones of Total Rare Earth Oxides including Yttrium ("**TREO**") intercepts from surface including **105m at 6.97% TREO** which is mineralised to end of hole and includes **35m at 8.50% TREO**
- **105m at 7.17% TREO** from surface to end of hole including **53m at 8.12% TREO**
- **100m at 6.18% TREO** from surface to end of hole including **13m at 10.59% TREO**; and
- **86m at 6.23% TREO** from surface.

Niobium

- **105m at 0.84% Nb₂O₅** from surface to end of hole including **31m at 1.15% Nb₂O₅** from 5m and **15m @ 1.11% Nb₂O₅** from 41m; and
- **105m at 0.66% Nb₂O₅** from surface to end of hole including **37m at 0.98% Nb₂O₅** from 9m and **5m @ 1.43% Nb₂O₅** from 100m to end of hole.

President/CEO Comments

PAW's President and CEO, Mr. Darren Townsend, commented, "With the receipt of these further seven high grade holes we can now finalise the updated Niobium Resource Estimate and Inaugural Rare Earth Resource for Mrima Hill which we expect to complete within the next four weeks."

Background to Drill Program and Results

During 2012 a total of 51 RC holes for 4,941m were drilled at Mrima Hill. Of this 37 holes for 3,482 metres

had not been sent for analysis. The 7 RC holes forming the basis of this News Release are situated across the center of the southern zone of the high grade TREO area identified by the Kenyan Mines Department in the 1950's (refer to contours on **Figure 1**).

These results are from holes drilled vertically between 54-105m depth. All holes reported with the exception of Holes PAWRC068 end in mineralisation, remaining open at depth in highly prospective clays.

Figure 1 shows a plan location of the RC drill holes as drilled

Figure 2 depicts a plan view of significant results for TREO mineralization

Figures 3 and 4 show cross sections through the TREO mineralization

To view Figures 1-4, please click on the following link: http://media3.marketwire.com/docs/paw715_F1-4.pdf.

Significant Rare Earth Results

DRILL HOLE INTERSECTION

PAWRC063: 105m at 6.97% TREO from surface to end of hole including 35m at 8.50% TREO from 43m.

PAWRC068: 67m at 5.59% TREO from surface and;
8m at 1.73% TREO from 69m and;
3m at 1.24% TREO from 94m.

PAWRC071: 53m at 6.04% TREO from surface and;
37m at 2.16% TREO from 55m and;
8m at 6.87% TREO from 94m to end of hole.

PAWRC072: 105m at 7.17% TREO from surface to end of hole including 25m at 8.51% TREO from 13m and 53m at 8.12% TREO from 41m.

PAWRC075: 100m at 6.18% TREO from surface to end of hole including 13m at 10.59% TREO from surface and 61m at 6.65% TREO from 19m.

PAWRC076: 54m at 5.78% TREO from surface to end of hole including 35m at 7.87% TREO from surface.

PAWRC082: 86m at 6.23% TREO from surface to end of hole* including 36m at 6.99% TREO from 1m.

* - Hole depth is 87m however final sample is missing

Significant Niobium Results

DRILL HOLE INTERSECTION

PAWRC063: 105m at 0.66% Nb₂O₅ from surface to end of hole including 37m at 0.98% Nb₂O₅ from 9m and 5m at 1.43% Nb₂O₅ from 100m.

PAWRC068: 6m at 0.42% Nb₂O₅ from surface and;
24m at 0.90% Nb₂O₅ from 13m and;
60m at 0.74% Nb₂O₅ from 42m.

PAWRC071: 13m at 0.44% Nb₂O₅ from surface and;
43m at 0.76% Nb₂O₅ from 17m including 6m at 2.15% Nb₂O₅ from 39m.

PAWRC072: 105m at 0.84% Nb₂O₅ from surface to end of hole including 31m at 1.15% Nb₂O₅ from 5m and 15m at 1.11% Nb₂O₅ from 41m.

PAWRC075: 81m at 0.45% Nb₂O₅ from surface.

PAWRC076: 38m at 0.70% Nb₂O₅ from surface.

PAWRC082: 77m at 0.42% Nb₂O₅ from surface including 16m at 0.70% Nb₂O₅ from 32m.

All samples collected were from the cyclone on the RC drill rig with samples fed through a sample splitter. A sample size of 1-3 kilograms was then submitted to the Nagrom assay labs in Perth with sample preparation being undertaken by the Nagrom assay laboratory. Prepared samples were then dispatched to both the Ultratrace (Nb₂O₅ analysis by XRF) and Genalysis (for rare earth and matrix analysis using peroxide fusion and acid dissolution by ICP-MS) assay labs in Perth, Western Australia.

Table 1 provides a summary of the TREO results from the Genalysis laboratory in Perth and **Table 2** has a summary of Niobium results from the Ultratrace laboratory in Perth.

To view Tables 1 and 2, please click on the following link:

