

Osisko Infill Drilling Continues to Intersect High-Grade at Lynx

08.07.2019 | [GlobeNewswire](#)

TORONTO, July 08, 2019 - [Osisko Mining Inc.](#) (OSK:TSX. "Osisko" or the "Corporation") is pleased to provide new infill drilling results from the ongoing definition and expansion program at its 100% owned Windfall gold project located in the Abitibi greenstone belt, Urban Township, Eeyou Istchee James Bay, Québec. The program is currently focused on infill drilling within the main Windfall gold deposit and the adjacent Lynx deposit (located immediately NE of Windfall), exploration and expansion drilling on the main mineralized zones, and deep exploration in the central areas of the intrusive system.

Significant new analytical results from 35 intercepts in 10 drill holes and 13 wedges from surface focused on Lynx and Main Zone infill drilling are presented below. Additionally, 34 intercepts in 20 underground drill holes focused on 5 metres infill drilling, including the pending Lynx bulk sample area, are included in a second table below.

Highlights from new infill drilling results at Lynx include: 84.6 g/t Au over 11.8 metres in OSK-W-19-909-W11; 429 g/t Au over 2.2 metres in OSK-W-19-934-W2; 149 g/t Au over 2.2 metres in WST-19-0134; 121 g/t Au over 2.3 metres in OSK-W-19-1857-W2; 119 g/t Au over 2.0 metres in WST-19-0139; 86.9 g/t Au over 2.7 metres in WST-19-0145 and 63.2 g/t Au over 2.3 metres in WST-19-0138. Maps showing hole locations and full analytical results are available at www.osiskomining.com.

Osisko President and Chief Executive Officer John Burzynski commented: "We continue to be impressed by the gold mineralization we are encountering in the infill and expansion drilling at Lynx. We are very optimistic that we will be able to add significant ounces as we progress with resource drilling and exploration of the Windfall deposit."

Surface Drilling

Hole No.	From (m)	To (m)	Interval (m)	Au (g/t) uncut	Au (g/t) cut to 100 g/t Area	Zone
OSK-W-18-1539	1021.8	1024.0	2.2	4.46		Lynx 4
<i>including</i>	1021.8	1022.5	0.7	13.2		Lynx 4
OSK-W-18-1745	800.9	803.4	2.5	5.96		Lynx 4
<i>including</i>	801.6	802.1	0.5	29.2		Lynx 4
	905.9	911.6	5.7	3.98		Lynx_327
OSK-W-19-909-W7	985.0	987.1	2.1	14.3		Lynx_313
OSK-W-19-909-W11	912.2	924.0	11.8	84.6	42.3	
<i>including</i>	915.8	922.6	6.8	142	68.2	
<i>including</i>	915.8	917.4	1.6	149	100	Lynx_317
<i>including</i>	920.1	922.6	2.5	266	97.2	
OSK-W-19-934-W2	831.0	834.0	3.0	32.9		Lynx 4
<i>including</i>	831.0	832.0	1.0	79.8		Lynx 4
	855.9	857.9	2.0	16.9		Lynx 4
<i>including</i>	856.9	857.9	1.0	33.4		Lynx 4
	901.8	904.0	2.2	429	50.2	
<i>including</i>	901.8	902.4	0.6	1270	100	Lynx_313
<i>including</i>	903.5	904.0	0.5	361	100	
OSK-W-19-1181-W8	910.3	912.6	2.3	8.31		Lynx_314
<i>including</i>	911.0	911.6	0.6	28.3		Lynx_314

	933.7	936.0	2.3	13.1			
<i>including</i>	934.7	935.3	0.6	34.2		Lynx_317	Lynx
OSK-W-19-1181-W9	952.4	955.0	2.6	48.2	28.3		
<i>including</i>	953.4	954.1	0.7	174	100	Lynx_317	Lynx
OSK-W-19-1746-W1	683.2	685.7	2.5	19.0			
<i>including</i>	683.2	683.7	0.5	50.1		Lynx	Lynx
	764.0	766.4	2.4	17.0			
<i>including</i>	765.2	765.7	0.5	81.4		Lynx_336	Lynx
OSK-W-19-1835	946.0	948.4	2.4	5.06		Vein	
OSK-W-19-1835-W1	651.2	653.3	2.1	4.28			
<i>including</i>	651.2	652.4	1.2	7.38		Underdog	Underdog
	862.8	865.0	2.2	3.26		Underdog	Underdog
OSK-W-19-1857-W2	1274.5	1276.8	2.3	121	39.7		
<i>including</i>	1274.9	1275.8	0.9	307	100	Lynx_313	Lynx
OSK-W-19-1875	244.3	247.0	2.7	5.76			
<i>including</i>	244.3	245.1	0.8	18.3		Vein	
OSK-W-19-1880	149.0	151.2	2.2	4.07			
<i>including</i>	149.6	150.2	0.6	14.8		Caribou corridor	Caribou
OSK-W-19-1900	262.0	264.0	2.0	9.41			
<i>including</i>	262.0	263.0	1.0	18.3		Caribou_230	Caribou
OSK-W-19-1921-W1	787.0	789.1	2.1	19.9			
<i>including</i>	788.3	788.8	0.5	81.3		Lynx	Lynx
	897.6	900.0	2.4	4.01		Lynx_312	Lynx
OSK-W-19-1921-W2	1040.2	1043.7	3.5	7.83		Lynx 4	Lynx
OSK-W-19-1932	671.7	675.8	4.1	31.3			
<i>including</i>	674.1	675.0	0.9	97.3		Lynx_326	Lynx
OSK-W-19-1932-W1	688.3	690.5	2.2	3.40		Lynx_324	Lynx
OSK-W-19-1932-W2	717.5	719.5	2.0	10.2		Lynx_326	Lynx
	745.0	747.0	2.0	86.5	70.4		
<i>including</i>	745.6	747.0	1.4	123	100	Lynx_324	Lynx
	800.9	803.0	2.1	85.8	20		
<i>including</i>	801.8	802.2	0.4	445	100	Lynx 4	Lynx
	808.0	810.0	2.0	29.6	27.2		
<i>including</i>	808.5	809.0	0.5	110	100	Lynx 4	Lynx
OSK-W-19-1939	351.0	353.0	2.0	6.44		Lynx_316	Lynx
	357.9	360.3	2.4	18.7			
<i>including</i>	357.9	358.4	0.5	68.3		Lynx_316	Lynx
OSK-W-19-1942	861.2	863.5	2.3	11.3			
<i>including</i>	861.2	861.7	0.5	45.4		Lynx_327	Lynx
OSK-W-19-1942-W1	837.0	839.0	2.0	3.08		Lynx_313	Lynx
OSK-W-19-1949	731.0	733.0	2.0	3.19		Lynx	Lynx
	971.3	975.0	3.7	11.5			
<i>including</i>	972.1	973.1	1.0	31.2		Lynx_317	Lynx
	996.2	999.5	3.3	31.3			
<i>including</i>	996.2	997.1	0.9	74.2		Lynx_330	Lynx
<i>including</i>	999.1	999.5	0.4	79.0			

Underground drilling

Hole No.	From (m)	To (m)	Interval (m)	Au (g/t) uncut	Au (g/t) cut to 100 g/t	Area	Zone
WST-19-0088	58.9	61.0	2.1	4.67		Caribou_201	Caribou
WST-19-0112	70.8	73.1	2.3	17.2		Lynx_311	Lynx

WST-19-0119	108.0	110.0	2.0	3.07		Lynx_305	Lynx
WST-19-0123	113.6	116.0	2.4	6.77		Lynx_304	Lynx
WST-19-0127B	41.3	44.4	3.1	21.9	13.8	Lynx_311	Lynx
<i>including</i>	41.9	42.2	0.3	185	100		
WST-19-0131	50.0	52.4	2.4	13.8		Lynx_311	Lynx
<i>including</i>	50.0	50.3	0.3	75.2			
	56.6	59.1	2.5	92.1	28.6		
<i>including</i>	58.0	58.3	0.3	393	100	Lynx_308	Lynx
<i>including</i>	58.7	59.1	0.4	277	100		
	82.5	84.6	2.1	12.9		Lynx_305	Lynx
WST-19-0133	52.9	59.1	6.2	12.9	12.4		
<i>including</i>	52.9	53.2	0.3	75.4		Lynx_311	Lynx
<i>including</i>	57.0	57.3	0.3	42.7			
<i>including</i>	58.3	58.6	0.3	109	100		
WST-19-0134	51.9	54.0	2.1	10.8		Lynx_308	Lynx
<i>including</i>	51.9	52.2	0.3	68.4			
	71.5	73.7	2.2	10.5		Lynx_310	Lynx
<i>including</i>	72.2	72.8	0.6	38.2			
	110.8	113.0	2.2	149	27.3	Lynx_304	Lynx
<i>including</i>	110.8	111.1	0.3	994	100		
WST-19-0137	96.0	98.0	2.0	10.4		Lynx_305	Lynx
WST-19-0138	68.0	70.4	2.4	5.52		Lynx	Lynx
<i>including</i>	68.5	69.4	0.9	14.5			
	96.0	98.0	2.0	3.04		Lynx_304	Lynx
<i>including</i>	96.5	97.1	0.6	9.91			
	100.0	102.3	2.3	63.2	39.3	Lynx_304	Lynx
<i>including</i>	100.6	101.5	0.9	161	100		
WST-19-0139	103.0	105.0	2.0	119	46.6	Lynx_304	Lynx
<i>including</i>	103.0	103.9	0.9	262	100		
WST-19-0140	42.0	44.0	2.0	7.58		Lynx	Lynx
<i>including</i>	43.1	43.4	0.3	48.2			
	53.4	57.7	4.3	9.00		Lynx_308	Lynx
<i>including</i>	53.4	53.8	0.4	76.0			
	102.8	105.4	2.6	10.3		Lynx_304	Lynx
WST-19-0145	68.8	71.5	2.7	86.9	15.2	Lynx_310	Lynx
<i>including</i>	68.8	69.1	0.3	746	100		
WST-19-0146	63.0	65.0	2.0	4.84		Lynx_308	Lynx
<i>including</i>	63.8	64.1	0.3	31.8			
	77.0	79.0	2.0	3.51		Lynx_310	Lynx
	88.0	90.0	2.0	63.7	50.2	Lynx_310	Lynx
<i>including</i>	89.0	90.0	1.0	127	100		
	105.4	109.4	4.0	5.60		Lynx_305	Lynx
WST-19-0149	49.0	51.2	2.2	106	68.4	Lynx_311	Lynx
<i>including</i>	49.0	49.4	0.4	199	100		
<i>including</i>	50.7	51.2	0.5	188	100		
WST-19-0150B	44.0	46.1	2.1	17.7		Lynx_311	Lynx
	49.8	51.9	2.1	33.5	26.7	Lynx_311	Lynx
<i>including</i>	49.8	50.2	0.4	136	100		
WST-19-0153	45.9	50.4	4.5	15.9		Lynx_308	Lynx
<i>including</i>	45.9	47.0	1.1	51.0			
WST-19-0156	46.0	48.9	2.9	10.4		Lynx_311	Lynx
<i>including</i>	47.8	48.6	0.8	30.2			

	98.5	101.0	2.5	21.3	16.5	Lynx_305	Lynx
<i>including</i>	99.6	100.0	0.4	130	100		
WST-19-0157	70.0	72.0	2.0	3.86		Lynx_308	Lynx
WST-19-0158	119.3	121.3	2.0	10.3		Lynx_305	Lynx
	132.0	134.0	2.0	3.10		Lynx_304	Lynx

Notes: True widths are estimated at 55 – 80% of the reported core length interval. See "Quality Control and Reporting Protocols" below.

Surface Drilling

Hole Number	Azimuth (°)	Dip (°)	Length (m)	UTM E	UTM N	Elevation (m)	Section
OSK-W-18-1539	145	-52	1119	453374	5435448	402	3750
OSK-W-18-1745	134	-51	1038	453218	5435347	407	3575
OSK-W-19-909-W7	131	-55	1080	453683	5435677	401	4125
OSK-W-19-909-W11	131	-55	990	453683	5435677	401	4125
OSK-W-19-934-W2	144	-55	1167	453407	5435463	401	3800
OSK-W-19-1181-W8	133	-58	1035	453789	5435790	401	4275
OSK-W-19-1181-W9	133	-58	1058	453789	5435790	401	4275
OSK-W-19-1746-W1	142	-53	840	453280	5435389	405	3650
OSK-W-19-1835	172	-53	1293	452305	5435389	406	2825
OSK-W-19-1835-W1	172	-53	948	452305	5435474	406	2825
OSK-W-19-1857-W2	108	-58	1449	453525	5435704	405	4000
OSK-W-19-1875	154	-46	360	452021	5435223	405	2450
OSK-W-19-1880	333	-52	372	452553	5434682	403	2650
OSK-W-19-1900	323	-52	345	452619	5434731	402	2750
OSK-W-19-1921-W1	112	-52	951	453501	5435490	399	3875
OSK-W-19-1921-W2	112	-52	1080	453501	5435490	399	3875
OSK-W-19-1932	127	-55	711	453709	5435621	400	4125
OSK-W-19-1932-W1	127	-55	723	453709	5435621	400	4125
OSK-W-19-1932-W2	127	-55	824	453709	5435621	400	4125
OSK-W-19-1939	126	-45	519	453285	5435277	400	3600
OSK-W-19-1942	128	-54	930	453314	5435388	403	3675
OSK-W-19-1942-W1	128	-54	942	453314	5435388	403	3675
OSK-W-19-1949	105	-57	1073	453440	5435479	401	3825

Underground drilling

Hole Number	Azimuth (°)	Dip (°)	Length (m)	UTM E	UTM N	Elevation (m)	Section
WST-19-0088	307	16	136	452147	5434645	194	2300
WST-19-0112	141	-28	139	453219	5435116	223	3450
WST-19-0119	144	-35	157	453252	5435110	206	3475
WST-19-0123	132	-17	135	453253	5435110	206	3475
WST-19-0127B	154	-5	55	453252	5435110	206	3475
WST-19-0131	143	24	129	453252	5435110	206	3475
WST-19-0133	154	27	129	453252	5435110	206	3475
WST-19-0134	154	14	127	453251	5435110	207	3475
WST-19-0137	136	13	129	453268	5435108	206	3500
WST-19-0138	136	3	120	453268	5435108	206	3500
WST-19-0139	134	-8	121	453266	5435107	206	3500
WST-19-0140	134	-16	120	453266	5435107	206	3500

WST-19-0145	142	-45	142	453290	5435116	204	3525
WST-19-0146	175	-44	160	453289	5435116	204	3525
WST-19-0149	159	-28	90	453252	5435110	205	3475
WST-19-0150B	145	-14	72	453252	5435110	205	3475
WST-19-0153	162	25	115	453289	5435116	206	3525
WST-19-0156	161	-39	142	453289	5435116	205	3525
WST-19-0157	164	-47	103	453289	5435116	205	3525
WST-19-0158	149	-50	151	453289	5435116	204	3525

OSK-W-18-1539 intersected 4.46 g/t Au over 2.2 metres. Mineralization includes up to 10% disseminated pyrite, 4% pyrite stringers and clusters, 3% pyrite in ptymatic tourmaline veinlets and 2% interstitial pyrite within a silica, sericite and fuchsite altered rhyolite and a sericite and chlorite altered andesite.

OSK-W-18-1745 intersected two intervals: 5.96 g/t Au over 2.5 metres and 3.98 g/t Au over 5.7 metres. The first interval contains 7% disseminated and dismembered stringer pyrite in a moderate silica and sericite altered andesite at the contact with a porphyritic felsic dike. The second interval contains up to 15% disseminated, clustered, and stringer pyrite in quartz-tourmaline and quartz-carbonate veins at the contact between a sericite and silica altered rhyolite and a sericite, silica and fuchsite altered gabbro.

OSK-W-19-909-W7 intersected 14.3 g/t Au over 2.1 metres. Mineralization includes 1% pyrite stringers in a strong silica, moderate sericite and weak fuchsite altered porphyritic felsic dike.

OSK-W-19-909-W11 intersected 84.6 g/t Au over 11.8 metres. Mineralization includes local visible gold, up to 15% pyrite stringers and clusters, and trace chalcopyrite and sphalerite within a strongly sericitized and silicified gabbro with moderate fuchsite alteration.

OSK-W-19-934-W2 intersected three intervals: 32.9 g/t Au over 3.0 metres, 16.9 g/t Au over 2.0 metres and 429 g/t Au over 2.2 metres. The first interval contains 4% pyrite stringers and clusters hosted in a moderate sericite-silica altered and bleached rhyolite. The second interval contains 5% interstitial pyrite in a moderate chlorite-silica altered rhyolite with breccia texture. The third interval contains 1% local visible gold veinlets and clusters and 5% pyrite within a smoky quartz-tourmaline crustiform vein hosted in a strong sericite and silica altered rhyolite at the contact with a bleached andesite.

OSK-W-19-1181-W8 intersected 8.31 g/t Au over 2.3 metres and 13.1 g/t Au over 2.3 metres. The first interval contains 5% disseminated, stringer, and clustered pyrite, and trace chalcopyrite in quartz-carbonate veinlets with local visible gold within a strong silica and moderate sericite altered gabbro. The second interval contains up to 30% disseminated, clustered and locally semi-massive pyrite in a strong silica altered rhyolite with moderate sericite alteration at the contact with a mafic volcanic unit.

OSK-W-19-1181-W9 intersected 48.2 g/t Au over 2.6 metres. Mineralization includes up to 5% disseminated pyrite, 3% pyrite clusters and trace chalcopyrite with pervasive-silica flooding and local sericite alteration hosted in a gabbro at the contact with a sericitized rhyolite.

OSK-W-19-1746-W1 intersected 19.0 g/t Au over 2.5 metres and 17.0 g/t Au over 2.4 metres. The first interval contains local visible gold with 20% pyrite filling millimeter to centimeter scale tourmaline fractures with massive pervasive silica flooding hosted in a porphyritic felsic intrusion. The second interval contains local visible gold with 15% disseminated or fracture-filling pyrite and 1% sphalerite in a smoky quartz vein. Both intervals are hosted in a silicified rhyolite with local fragmental texture.

OSK-W-19-1835 intersected 5.06 g/t Au over 2.4 metres. Mineralization includes 1% disseminated pyrite with quartz veins in a weak sericite altered porphyritic felsic dike.

OSK-W-19-1835-W1 intersected 4.28 g/t Au over 2.1 metres and 3.26 g/t Au over 2.2 metres. The first interval contains 2% pyrite stringers, 10% semi-massive pyrite within a sericite and silica altered porphyritic felsic dike. The second interval contains 3% pyrite within pervasive silica flooding, 1% fracture filling pyrite and 1% disseminated pyrite in a bleached porphyritic felsic dike.

OSK-W-19-1857-W2 intersected 121 g/t Au over 2.3 metres. Mineralization includes local visible gold in quartz-carbonate veins, 10% disseminated, clustered, and stringer pyrite ± tourmaline, and trace chalcopyrite within a rhyolite with strong silica and moderate sericite alteration.

OSK-W-19-1875 intersected 5.76 g/t Au over 2.7 metres. Mineralization includes 6% disseminated pyrite and 1% pyrite in quartz-carbonate veins hosted in a sericitized andesite.

OSK-W-19-1880 intersected 4.07 g/t Au over 2.2 metres. Mineralization includes 3% pyrite-tourmaline stringers and trace disseminated pyrite with pervasive silica flooding and tourmaline pygmatic veins at the contact between a porphyritic felsic dike, a rhyolite and a gabbro.

OSK-W-19-1900 intersected 9.41 g/t Au over 2.0 metres. Mineralization includes 3% pyrite stringers with local silicification hosted in a weak to moderate sericite and silica altered porphyritic felsic dike.

OSK-W-19-1921-W1 intersected two intervals in Lynx: 19.9 g/t Au over 2.1 metres and 4.01 g/t Au over 2.4 metres. The first interval contains local visible gold with 5% disseminated and stringer pyrite in a smoky quartz vein hosted in a gabbro with local and strong pervasive silica flooding. The second interval contains 4% pyrite stringers, 2% pyrite clusters and 20% quartz-carbonate veins in a moderate chlorite altered gabbro with weak sericite, silica and fuchsite alteration.

OSK-W-19-1921-W2 intersected 7.83 g/t Au over 3.5 metres. Mineralization includes 3% pyrite-tourmaline stringers and trace disseminated pyrite with quartz-tourmaline veins in a weakly sericitized and silicified rhyolite.

OSK-W-19-1932 intersected 31.3 g/t Au over 4.1 metres. Mineralization includes 5% pyrite clusters and 2% pyrite stringers within a local quartz-carbonate vein hosted in a weak fuchsite, moderate chlorite and sericite altered gabbro.

OSK-W-19-1932-W1 intersected 3.40 g/t Au over 2.2 metres. Mineralization includes 5% pyrite clusters and 1% pyrite stringers at the contact between a weakly sericitized and moderately silicified rhyolite and a weak silica-chlorite and moderate sericite altered gabbro.

OSK-W-19-1932-W2 intersected four intervals: 10.2 g/t Au over 2.0 metres, 86.5 g/t Au over 2.0 metres, 85.8 g/t Au over 2.1 metres and 29.6 g/t Au over 2.0 metres. The first interval contains 4% pyrite stringers and clusters in a sericitized and silicified gabbro with moderate fuchsite alteration. The second interval contains 5% pyrite clusters and stringers in a fragmental felsic unit with strong local silica and fuchsite alteration. The third and fourth intervals contain local visible gold and 3% pyrite with pervasive silica flooding within a sericitized and silicified rhyolite.

OSK-W-19-1939 intersected 6.44 g/t Au over 2.0 metres and 18.7 g/t Au over 2.4 metres. Mineralization in both intervals contains local visible gold, 5% disseminated, stringer, and clustered pyrite with pygmatic tourmaline veinlets hosted in a moderately sericitized and weakly silicified rhyolite.

OSK-W-19-1942 intersected 11.3 g/t Au over 2.3 metres. Mineralization includes local visible gold, 2% disseminated and stringer pyrite with local tourmaline with pervasive silica flooding hosted in a moderate sericite and weak silica altered rhyolite.

OSK-W-19-1942-W1 intersected 3.08 g/t Au over 2.0 metres. Mineralization includes 4% pyrite stringers and clusters and trace sphalerite and chalcopyrite hosted in a moderate sericite and silica altered rhyolite at the contact with a moderate chlorite altered gabbro.

OSK-W-19-1949 intersected three intervals: 3.19 g/t Au over 2.0 metres, 11.5 g/t Au over 3.7 metres and 31.3 g/t Au over 3.3 metres. The first interval contains local visible gold and smoky quartz veins with up to 8% pyrite in a porphyritic felsic dike with weak fuchsite and carbonate alteration. The second interval contains 4% pyrite stringers and clusters, 2% interstitial sphalerite and trace chalcopyrite with pervasive

silica flooding, quartz carbonate veins and pygmatic tourmaline veins hosted in a silicified felsic intrusion at the contact with a sericitized rhyolite. The third interval contains up to 10% pyrite with brecciated smoky quartz veins hosted in a moderate sericite and weak silica altered rhyolite.

WST-19-0088 intersected 4.67g/t Au over 2.1 metres. Mineralization includes 3% pyrite clusters and stringers and 1% disseminated pyrite within a chloritized fragmental andesite unit. WST-19-0088 was drilled from underground drill station AN-190-155-N located 190 metres below surface from section 2300E.

WST-19-0112 intersected 17.2 g/t Au over 2.3 metres. Mineralization includes 7% clustered and disseminated pyrite with intense pervasive silica flooding in a silicified porphyritic felsic dike and a rhyolite. WST-19-0112 was drilled from underground drill station RA-180-190-E located 180 metres below surface from section 3450E.

WST-19-0119 intersected 3.07 g/t Au over 2.0 metres. Mineralization includes 2% pyrite clusters, trace disseminated pyrite and 10% quartz-tourmaline in a weak silica and sericite altered rhyolite. WST-19-0119 was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0123 intersected 6.77 g/t Au over 2.0 metres. Mineralization includes 3% pyrite clusters with pervasive silica flooding within a weak sericite altered rhyolite. WST-19-0123 was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0127B intersected 21.9 g/t Au over 3.1 metres. Mineralization includes 3% clustered and disseminated pyrite, local pyrite-tourmaline stringers and trace sphalerite with smoky pervasive silica flooding and local pyritized fragments. Local visible gold is observed with pyrite within a sericitized breccia hosted in a sericitized rhyolite. WST-19-0127B was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0131 intersected three intervals: 13.8 g/t Au over 2.4 metres, 92.1 g/t Au over 2.5 metres and 12.9 g/t Au over 2.1 metres. Mineralization in all three intervals contains local visible gold with 2% pyrite in pervasive silica flooding zone and trace sphalerite in a sericitized and silicified fragmental felsic unit. WST-19-0131 was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0133 intersected 12.9 g/t Au over 6.2 metres. Mineralization includes local visible gold, 1% pyrite clusters with 2% pervasive silica flooding. Mineralization is directly related to silica fracture filling. The interval is hosted in a sericitized fragmental felsic unit. WST-19-0133 was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0134 intersected three intervals: 10.8 g/t Au over 2.1 metres, 10.5 g/t Au over 2.2 metres and 149 g/t Au over 2.2 metres. The first and second intervals contain local visible gold, 2% disseminated, clustered, and stringer pyrite with pervasive silica flooding hosted in a sericitized fragmental felsic unit. The third interval contains local visible gold, 7% pyrite stringers and trace sphalerite with pervasive silica flooding hosted in a weakly sericitized and moderately silicified porphyritic rhyolite. WST-19-0134 was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0137 intersected 10.4 g/t Au over 2.0 metres. Mineralization includes trace disseminated pyrite in quartz-carbonate veins in a moderated chlorite altered and faulted rhyolite. WST-19-0137 was drilled from underground drill station AN-195-265-S located 195 metres below surface from section 3500E.

WST-19-0138 intersected three intervals: 5.52 g/t Au over 2.4 metres, 3.04 g/t Au over 2.0 metres and 63.2 g/t Au over 2.3 metres. The first interval contains trace pyrite with quartz-tourmaline veins in a moderate chlorite-sericite altered and fragmental felsic unit. The second interval contains 4% pyrite clusters with two intense pervasive silica flooding zones with fragments of felsic intrusion, volcanics, tourmaline and pyrite at the contact with a sericitized rhyolite. The third interval contains traces of disseminated and clustered pyrite with 30% quartz-carbonate veins at the contact between a rhyolite and a chloritized gabbro. WST-19-0138 was drilled from underground drill station AN-195-265-S located 195 metres below surface from section 3500E.

WST-19-0139 intersected 119 g/t Au over 2.0 metres. Mineralization includes local visible gold and 15% pyrite within smoky quartz vein or pervasive silica flooding hosted at the strongly silicified contact between a felsic porphyritic intrusion and a rhyolite. WST-19-0139 was drilled from underground drill station AN-195-265-S located 195 metres below surface from section 3500E.

WST-19-0140 intersected three interval: 7.58 g/t Au over 2.0 metres, 9.00 g/t Au over 4.3 metres and 10.3 g/t Au over 2.6 metres. The first interval contains trace pyrite stringers, 2% disseminated pyrite and trace sphalerite with irregular quartz-carbonate-tourmaline veins hosted in sericitized fragmental felsic unit at the contact with a sericitized rhyolite. The second interval contains 1% sphalerite clusters and 7% pyrite within smoky veins hosted in weakly silicified, sericitized and fuchsitized rhyolite. The third interval contains 1% disseminated and clustered pyrite with intense silica flooding hosted in a silicified porphyritic felsic dike at the contact with a sericitized and silicified rhyolite. WST-19-0140 was drilled from underground drill station AN-195-265-S located 195 metres below surface from section 3500E.

WST-19-0145 intersected 86.9 g/t Au over 2.7 metres. Mineralization includes local visible gold, trace sphalerite and 1% disseminated pyrite with quartz-tourmaline-carbonate veins within a silica and carbonate altered rhyolite. WST-19-0145 was drilled from underground drill station BM-200-285-S located 200 metres below surface from section 3525E.

WST-19-0146 intersected four intervals: 4.84 g/t Au over 2.0 metres, 3.51 g/t Au over 2.0 metres, 63.7 g/t Au over 2.0 metres and 5.60 g/t Au over 4.0 metres. The first interval contains 5% semi-massive pyrite in a 10 centimetres-wide band, 1% pyrite clusters, and 1% pyrite-tourmaline stringers in a silica altered rhyolite. The second interval contains 30% pyrite with quartz-tourmaline veins in a sericitized rhyolite. The third interval contains trace disseminated pyrite in stockwork in a weakly sericitized rhyolite. The fourth interval contains trace pyrite in a breccia with tourmaline in a moderate chlorite, silica and sericite altered gabbro. WST-19-0146 was drilled from underground drill station BM-200-285-S located 200 metres below surface from section 3525E.

WST-19-0149 intersected 106 g/t Au over 2.2 metres. Mineralization includes local visible gold, 3% dismembered pyrite stringers, 2% pyrite clusters, trace sphalerite and chalcopyrite with weakly fuchsitized pervasive silica flooding hosted in a moderate sericite-silica altered rhyolite. WST-19-0149 was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0150B intersected 17.7g/t Au over 2.1 metres and 33.5 g/t Au over 2.1 metres in Lynx. Mineralization in both intervals contains local visible gold, 3% pyrite stringers and clusters and trace sphalerite with pervasive silica flooding and quartz-carbonate veinlets hosted in a sericite and silica altered rhyolite. WST-19-0150B was drilled from underground drill station RE-195-265-O located 195 metres below surface from section 3475E.

WST-19-0153 intersected 15.9 g/t Au over 4.5 metres. Mineralization includes up to 5% pyrite stringers hosted in a moderately silicified felsic intrusion with intrusive, volcanic, tourmaline and pyrite fragments. WST-19-0153 was drilled from underground drill station BM-200-285-S located 200 metres below surface from section 3525E.

WST-19-0156 intersected 10.4 g/t Au over 2.9 metres and 21.3 g/t Au over 2.5 metres. Mineralization includes up to 15% pyrite associated with pervasive silica flooding hosted in a strongly silicified rhyolite. WST-19-0156 was drilled from underground drill station BM-200-285-S located 200 metres below surface from section 3525E.

WST-19-0157 intersected 3.86 g/t Au over 2.0 metres. Mineralization includes 3% pyrite with pervasive silica flooding in a moderate sericite, silica and chlorite altered gabbro. WST-19-0157 was drilled from underground drill station BM-200-285-S located 200 metres below surface from section 3525E.

WST-19-0158 intersected 10.3 g/t Au over 2.0 metres and 3.10 g/t Au over 2.0 metres. The first interval contains 5% pyrite stringers and clusters with pervasive silica flooding within a moderate sericite, silica and fuchsite altered gabbro. The second interval contains 3% pyrite clusters at the contact between a rhyolite and a strongly sericitized, bleached and fuchsite altered gabbro. WST-19-0158 was drilled from underground drill station BM-200-285-S located 200 metres below surface from section 3525E.

Qualified Person

The scientific and technical content of this news release has been reviewed, prepared and approved by Mr. Louis Grenier, M.Sc.A., P.Geo. (OGQ 800), Project Manager of Osisko's Windfall Lake gold project, who is a "qualified person" as defined by National Instrument 43-101 – Standards of Disclosure for Mineral Projects ("NI 43-101").

Quality Control and Reporting Protocols

True width determination is estimated at 55-80% of the reported core length interval for the zone. Assays are uncut except where indicated. Intercepts occur within geological confines of major zones but have not been correlated to individual vein domains at this time. Reported intervals include minimum weighted averages of 3.0 g/t Au diluted over core lengths of at least 2.0 metres. All NQ core assays reported were obtained by either 1-kilogram screen fire assay or standard 50-gram fire-assaying-AA finish or gravimetric finish at (i) ALS Laboratories in Val d'Or, Québec, Thunder Bay, Ontario, Sudbury, Ontario or Vancouver, British Columbia, or (ii) Bureau Veritas in Timmins, Ontario. The 1-kilogram screen assay method is selected by the geologist when samples contain coarse gold or present a higher percentage of pyrite than surrounding intervals. Selected samples are also analyzed for multi-elements, including silver, using an Aqua Regia-ICP-AES method at ALS Laboratories. Drill program design, Quality Assurance/Quality Control ("QA/QC") and interpretation of results is performed by qualified persons employing a QA/QC program consistent with NI 43-101 and industry best practices. Standards and blanks are included with every 20 samples for QA/QC purposes by the Corporation as well as the lab. Approximately 5% of sample pulps are sent to secondary laboratories for check assay.

About the Windfall Lake Gold Deposit

The Windfall Lake gold deposit is located between Val-d'Or and Chibougamau in the Abitibi region of Québec, Canada. The mineral resource defined by Osisko, as disclosed in the Windfall Lake Technical Report (as defined below) and November 27, 2018 Lynx resource update, comprises 2,874,000 tonnes at 8.17 g/t Au (754,000 ounces) in the indicated mineral resource category and 10,352,000 tonnes at 7.11 g/t Au (2,366,000 ounces) in the inferred mineral resource category. For details regarding the key assumptions, parameters and methods used to estimate the mineral resources presented in respect of the Windfall Lake gold project, please see the technical report entitled "Technical Report and Mineral Resource Estimate for the Windfall Lake Project, Windfall Lake and Urban-Barry Properties" and dated June 12, 2018 (effective date of May 14, 2018), which has been prepared by InnovExplo Inc. from Val-d'Or, Québec (the "Windfall Lake Technical Report") and the press release "Osisko Releases Mineral Resource Update for Lynx" dated November 27, 2018, which has been prepared by Osisko and reviewed and approved by Micon International, Ltd. from Toronto, Ontario. The Windfall Lake Technical Report and press release are available on Osisko's website at www.osiskomining.com and on SEDAR under Osisko's issuer profile at www.sedar.com. The Windfall Lake gold deposit is currently one of the highest-grade resource-stage gold projects in Canada. Mineralization occurs in four principal zones: Lynx, Zone 27, Caribou and Underdog. All zones comprise sub-vertical lenses following intrusive porphyry contacts plunging to the northeast. The deposit is well defined from surface to a depth of 900 metres and remains open along strike and at depth. Mineralization has been identified 30 metres from surface in some areas and as deep as 2,000 metres in others, with significant potential to extend mineralization down-plunge and at depth.

About Osisko Mining Inc.

Osisko is a mineral exploration company focused on the acquisition, exploration, and development of precious metal resource properties in Canada. Osisko holds a 100% interest in the high-grade Windfall Lake gold deposit located between Val-d'Or and Chibougamau in Québec and holds a 100% undivided interest in a large area of claims in the surrounding the Urban Barry area and nearby Quévillon area (over 2,700 square kilometres).

Cautionary Note Regarding Forward-Looking Information

This news release contains "forward-looking information" within the meaning of the applicable Canadian securities legislation that is based on expectations, estimates, projections and interpretations as at the date of this news release. The information in this news release about the Windfall Lake gold deposit being one of the highest grade resource-stage gold projects in Canada; the significance of results from the new infill drilling and ongoing drill definition and expansion program at the Windfall Lake gold project; the significance of assay results presented in this news release; the deposit remaining open along strike and at depth; potential depth extensions of the mineralized zones down-plunge and at depth; the actual mineralization of local visible gold; the current 800,000 metre drill program; the type of drilling included in the drill program; potential mineralization; the potential to extend mineralization up and down-plunge and at depth at the Windfall Lake gold deposit; the ability to realize upon any mineralization in a manner that is economic; the ability to complete any proposed exploration activities and the results of such activities, including the

continuity or extension of any mineralization; and any other information herein that is not a historical fact may be "forward-looking information". Any statement that involves discussions with respect to predictions, expectations, interpretations, beliefs, plans, projections, objectives, assumptions, future events or performance (often but not always using phrases such as "expects", or "does not expect", "is expected", "interpreted", "management's view", "anticipates" or "does not anticipate", "plans", "budget", "scheduled", "forecasts", "estimates", "believes" or "intends" or variations of such words and phrases or stating that certain actions, events or results "may" or "could", "would", "might" or "will" be taken to occur or be achieved) are not statements of historical fact and may be forward-looking information and are intended to identify forward-looking information. This forward-looking information is based on reasonable assumptions and estimates of management of the Corporation at the time such assumptions and estimates were made, and involves known and unknown risks, uncertainties and other factors which may cause the actual results, performance or achievements of Osisko to be materially different from any future results, performance or achievements expressed or implied by such forward-looking information. Such factors include, among others, risks relating to the ability of exploration activities (including drill results) to accurately predict mineralization; errors in management's geological modelling; the ability of Osisko to complete further exploration activities, including drilling; property interests in the Windfall Lake gold project; the ability of the Corporation to obtain required approvals and complete transactions on terms announced; the results of exploration activities; risks relating to mining activities; the global economic climate; metal prices; dilution; environmental risks; and community and non-governmental actions. Although the forward-looking information contained in this news release is based upon what management believes, or believed at the time, to be reasonable assumptions. Osisko cannot assure shareholders and prospective purchasers of securities of the Corporation that actual results will be consistent with such forward-looking information, as there may be other factors that cause results not to be as anticipated, estimated or intended, and neither Osisko nor any other person assumes responsibility for the accuracy and completeness of any such forward-looking information, Osisko does not undertake, and assumes no obligation, to update or revise any such forward-looking statements or forward-looking information contained herein to reflect new events or circumstances, except as may be required by law.

For further information please contact:
John Burzynski
President and Chief Executive Officer
Telephone: (416) 363-8653

Dieser Artikel stammt von [Rohstoff-Welt.de](https://www.rohstoff-welt.de)

Die URL für diesen Artikel lautet:

<https://www.rohstoff-welt.de/news/329715--Osisko-Infill-Drilling-Continues-to-Intersect-High-Grade-at-Lynx.html>

Für den Inhalt des Beitrages ist allein der Autor verantwortlich bzw. die aufgeführte Quelle. Bild- oder Filmrechte liegen beim Autor/Quelle bzw. bei der vom ihm benannten Quelle. Bei Übersetzungen können Fehler nicht ausgeschlossen werden. Der vertretene Standpunkt eines Autors spiegelt generell nicht die Meinung des Webseiten-Betreibers wieder. Mittels der Veröffentlichung will dieser lediglich ein pluralistisches Meinungsbild darstellen. Direkte oder indirekte Aussagen in einem Beitrag stellen keinerlei Aufforderung zum Kauf-/Verkauf von Wertpapieren dar. Wir wehren uns gegen jede Form von Hass, Diskriminierung und Verletzung der Menschenwürde. Beachten Sie bitte auch unsere [AGB/Disclaimer!](#)

Die Reproduktion, Modifikation oder Verwendung der Inhalte ganz oder teilweise ohne schriftliche Genehmigung ist untersagt!
Alle Angaben ohne Gewähr! Copyright © by Rohstoff-Welt.de -1999-2026. Es gelten unsere [AGB](#) und [Datenschutzrichtlinien](#).